

On the $D^*D\pi$ and $\bar{B}^*\bar{B}\pi$ coupling constants in the effective quark model with chiral symmetry

A. N. Ivanov^{*†}, N.I. Troitskaya[†]

*Institut für Kernphysik, Technische Universität Wien,
Wiedner Hauptstr. 8-10, A-1040 Vienna, Austria*

Abstract

The coupling constants $g_{D^*D\pi}$ and $g_{\bar{B}^*\bar{B}\pi}$ of the $D^*D\pi$ and $\bar{B}^*\bar{B}\pi$ strong interactions are computed in the effective quark model with chiral symmetry incorporating Heavy quark effective theory (HQET) and Chiral perturbation theory at the quark level (CHPT)_q with linear realization of chiral $U(3) \times U(3)$ symmetry. We predict $g_{D^*D\pi} = 5.2$ and $g_{\bar{B}^*\bar{B}\pi} = 14.3$. The $1/M_c$ corrections to the constant $g_{D^*D\pi}$ are calculated and make up less than 1%. The contribution of $1/M_b$ corrections to the constant $g_{\bar{B}^*\bar{B}\pi}$ is much less.

PACS number(s): 13.30.Ce, 12.39.Ki, 14.20.Lq, 14.20.Mr

^{*}E-mail: ivanov@kph.tuwien.ac.at, Tel.: +43-1-58801-5598, Fax: +43-1-5864203

[†]Permanent Address: State Technical University, Department of Theoretical Physics, 195251 St. Petersburg, Russian Federation

1 Introduction

The investigation of strong interactions $D^*D\pi$ and $\bar{B}^*\bar{B}\pi$ runs parallel to the investigation of a dominance of vector D^* and $\bar{B}^*$ mesons in the semileptonic decays $D \rightarrow \pi\ell\nu_\ell$ and $\bar{B} \rightarrow \pi\ell\nu_\ell$ related to the strong transitions $D \rightarrow \pi$ and $\bar{B} \rightarrow \pi$, respectively, the form factors of which can be measured experimentally. In turn, the computation of the coupling constant $g_{D^*D\pi}$ of the $D^*D\pi$ interaction has a self-determined meaning, since the transitions $D^* \rightarrow D\pi$ do exist on-mass-shell of interacting mesons. The probabilities of the decays $B(D^{*+} \rightarrow D^0\pi^+) = (68.1 \pm 1.3)\%$, $B(D^{*+} \rightarrow D^+\pi^0) = (30.8 \pm 0.8)\%$ and $B(D^{*0} \rightarrow D^0\pi^0) = (63.6 \pm 2.8)\%$ have been measured recently by CLEO Collaboration [1].

The numerical values of $g_{D^*D\pi}$ and $g_{\bar{B}^*\bar{B}\pi}$ are distributed from $g_{D^*D\pi} = 7 \pm 2$, $g_{\bar{B}^*\bar{B}\pi} = 15 \pm 4$ [2] and $g_{D^*D\pi} = 6.3 \pm 1.9$, $g_{\bar{B}^*\bar{B}\pi} = 14 \pm 4$ [3] up to $g_{D^*D\pi} = 12.5 \pm 1.0$, $g_{\bar{B}^*\bar{B}\pi} = 29 \pm 3$ [4]. For the more complete list of references to the numerical values of the coupling constants $g_{D^*D\pi}$ and $g_{\bar{B}^*\bar{B}\pi}$ we relegate readers to Table 1 of Ref.[4]. Most of these results have been obtained within QCD sum rules approach [5].

Recently [6] in the effective quark model with chiral $U(3) \times U(3)$ symmetry incorporating Heavy quark effective theory (HQET) [7–9] and Chiral perturbation theory at the quark level (CHPT)_q with linear realization of chiral $U(3) \times U(3)$ symmetry [10] the coupling constant $g_{D^*D\pi}$ has been computed at leading order in large c -quark mass M_c expansion with the following value $g_{D^*D\pi} = g_{D^{*+}D^0\pi^+} = 5.2 \pm \Delta$. Here Δ is a theoretical uncertainty of our effective quark model which on the estimate of [6] makes up about 50%. However, for all cases of the application of HQET, supplemented by (CHPT)_q, to the description of strong interactions of heavy-light mesons [11–16] the resultant agreements between theoretical and experimental data turn out much better.

A possible substantial change of the $g_{D^*D\pi}$ coupling constant value in HQET, supplemented by (CHPT)_q, one can expect at next-to-leading order in large M_c expansion. Indeed, as has been shown in Ref.[15] such corrections to the form factors of semileptonic D -meson decays can make up more than 30%.

This letter is to compute $g_{D^*D\pi}$ in HQET, supplemented by (CHPT)_q, to next-to-leading order in large M_c expansion and to extend the obtained results on the computation of the coupling constant $g_{\bar{B}^*\bar{B}\pi}$ defining the $\bar{B}^*\bar{B}\pi$ strong interaction.

However, we have found that the contribution of $1/M_c$ corrections to $g_{D^*D\pi}$ makes up less than 1%. In turn $1/M_b$ corrections, where M_b is the mass of the b -quark, to $g_{\bar{B}^*\bar{B}\pi}$ is much less. The numerical value of $g_{\bar{B}^*\bar{B}\pi}$ calculated at leading order in large M_b expansion equals: $g_{\bar{B}^*\bar{B}\pi} = 14.3$. Thus, our results turn out to be compared reasonably well with the predictions of Refs.[2,3]. Therefore, in complete agreement with Refs.[2,3] we predict much less contributions of vector mesons D^* and $\bar{B}^*$ to the form factors of semileptonic decays $D \rightarrow \pi\ell\nu_\ell$ and $\bar{B} \rightarrow \pi\ell\nu_\ell$ than it can be expected due to Ref.[4].

2 Calculation of coupling constants at leading order in heavy quark mass expansion

The coupling constant $g_{D^*D\pi} = g_{D^{*+}D^0\pi^+}$ of the $D^{*+} \rightarrow D^0\pi^+$ decay is defined by the matrix element

$$\langle out; \pi^+(q) D^0(p) | D^*(Q); in \rangle = \quad (2.1)$$

$$= i(2\pi)^4 \delta^{(4)}(Q - p - q) \frac{M(D^{*+}(Q) \rightarrow D^0(p)\pi^+(q))}{(2E_{\pi^+}V 2E_{D^0}V 2E_{D^{*+}}V)^{1/2}}$$

with

$$M(D^{*+}(Q) \rightarrow D^0(p)\pi^+(q)) = g_{D^*D\pi} q \cdot e(Q), \quad (2.2)$$

where E_i ($i = \pi^+, D^0$ or D^{*+}) is the energy of the i -meson, and V is the normalization volume, then $e^\nu(Q)$ is the 4-vector of the D^{*+} -meson polarization. Note that by definition the coupling constant $g_{D^*D\pi}$ in Eq.(2.2) involves a factor 2 with respect to that defined in Ref.[1] (see Eq.(8) of Ref.[6]).

The computation of the matrix element $M(D^{*+}(Q) \rightarrow D^0(p)\pi^+(q))$ at leading order in large N , number of quark colours, and M_c expansion has been carried out in Ref.[6]. The result reads

$$M(D^{*+}(Q) \rightarrow D^0(p)\pi^+(q)) = \sqrt{2}g_{\pi^+} \frac{\sqrt{M_D M_{D^*}}}{\bar{v}'} \ell n\left(\frac{\bar{v}'}{4m}\right) q \cdot e(Q). \quad (2.3)$$

This gives one the coupling constant

$$g_{D^{*+}D^0\pi^+} = g_{D^*D\pi} = \frac{4\pi\sqrt{2}}{\sqrt{N}} \frac{\sqrt{M_D M_{D^*}}}{\bar{v}'} \ell n\left(\frac{\bar{v}'}{4m}\right) = 5.2, \quad (2.4)$$

where we have set $M_c = M_D = 1.86$ GeV and $M_{D^*} = 2.00$ GeV [1,2], and $g_{\pi^+} = 4\pi/\sqrt{N}$ [6,11–16] at $N = 3$, then $\bar{v}' = 4\Lambda = 2.66$ GeV [6,11–16] and Λ is the cut-off in Euclidean 3-momentum space connected with Λ_χ , the scale of spontaneous breaking of chiral symmetry (SB χ S) in (CHPT) $_q$, via the relation $\Lambda = \Lambda_\chi/\sqrt{2} = 0.67$ GeV at $\Lambda_\chi = 0.94$ GeV [10]. The cut-off Λ appears due to the computation of heavy-light constituent quark loops describing strong low-energy interactions in HQET and (CHPT) $_q$ at scales $p < \Lambda_\chi$.

The coupling constant $g_{\bar{B}^* \bar{B} \pi}$ can be obtained from Eq.(2.4) by the change $M_c \rightarrow M_b$, $M_D \rightarrow M_{\bar{B}}$ and $M_{D^*} \rightarrow M_{\bar{B}^*}$

$$g_{\bar{B}^* \bar{B} \pi} = \frac{4\pi\sqrt{2}}{\sqrt{N}} \frac{\sqrt{M_{\bar{B}} M_{\bar{B}^*}}}{\bar{v}'} \ell n\left(\frac{\bar{v}'}{4m}\right) = 14.3, \quad (2.5)$$

where we have set $M_{\bar{B}^*} = M_{\bar{B}} = M_b = 5.3$ GeV [6].

Our predictions for the coupling constants $g_{D^*D\pi} = 5.2$ and $g_{\bar{B}^* \bar{B} \pi} = 14.3$ agree with those given in Refs.[2,3].

3 Coupling constants to next-to-leading order in heavy quark mass expansion

Now we should proceed to the analysis of next-to-leading order corrections in heavy quark mass expansion. We suggest to start with the computation of the $1/M_c$ corrections to the coupling constant $g_{D^*D\pi}$. In our approach, where the amplitudes are described in terms of constituent quark loops with virtual momenta restricted by the SB χ S scale $\Lambda_\chi \ll M_c$, $1/M_c$ corrections can be computed by direct expansion of the c -quark Green function

in the momentum representation [15,16]. Since the virtual momentum k is restricted by the $\text{SB}\chi\text{S}$ scale $\Lambda_\chi \ll M_c$, we can expand the c -quark Green function in powers of k/M_c and, holding only the terms of order $O(1/M_c)$, obtain [15,16]

$$\begin{aligned} \frac{1}{M_c - \hat{k} - \hat{p}} &= -\left(\frac{1 + \hat{v}}{2}\right) \frac{1}{v \cdot k + i0} - \\ &- \frac{1}{2M_c} \left[\frac{\hat{k}}{v \cdot k + i0} - \left(\frac{1 + \hat{v}}{2}\right) \frac{k^2}{(v \cdot k + i0)^2} \right] + O\left(\frac{1}{M_c^2}\right), \end{aligned} \quad (3.1)$$

where, as usually, we have set $p^\mu = M_D v^\mu = M_c v^\mu$. Thus, the amplitude $M(D^{*+}(Q) \rightarrow D^0(p)\pi^+(q))$ can be given by

$$\begin{aligned} M(D^{*+}(Q) \rightarrow D^0(p)\pi^+(q)) &= \\ &= M^{(0)}(D^{*+}(Q) \rightarrow D^0(p)\pi^+(q)) + M^{(1)}(D^{*+}(Q) \rightarrow D^0(p) \end{aligned} \quad (3.2)$$

$$\pi^+(q)), \quad (3.3)$$

where the term $M^{(0)}(D^{*+}(Q) \rightarrow D^0(p)\pi^+(q))$ is given by Eq.(2.3), while $M^{(1)}(D^{*+}(Q) \rightarrow D^0(p)\pi^+(q))$ reads

$$\begin{aligned} M^{(1)}(D^{*+}(Q) \rightarrow D^0(p)\pi^+(q)) &= g_D g_{D^*} \frac{g_{\pi^+}}{\sqrt{2}} \frac{1}{2M_c} \frac{N}{16\pi^2} e^\mu(Q) \int \frac{d^4 k}{\pi^2 i} \\ &\text{tr} \left\{ \frac{1}{m - \hat{k} + \hat{q}} \gamma^5 \frac{1}{m - \hat{k}} \gamma^5 \left[\frac{\hat{k} \gamma_\mu}{v \cdot k + i0} - \left(\frac{1 + \hat{v}}{2}\right) \gamma_\mu \frac{k^2}{(v \cdot k + i0)^2} \right] \right\}. \end{aligned} \quad (3.4)$$

After the calculation of the trace over Dirac matrices we reduce the r.h.s. of Eq.(3.4) to the form

$$\begin{aligned} M^{(1)}(D^{*+}(Q) \rightarrow D^0(p)\pi^+(q)) &= g_D g_{D^*} \frac{g_{\pi^+}}{\sqrt{2}} \frac{1}{2M_c} \frac{N}{16\pi^2} e^\mu(Q) \times \\ &\left\{ \int \frac{d^4 k}{\pi^2 i} \frac{4(m^2 - k^2) k_\mu + 2 k^2 q_\mu}{[m^2 - (k - q)^2 - i0][m^2 - k^2 - i0]} \frac{1}{[v \cdot k + i0]} - \right. \\ &\left. - \int \frac{d^4 k}{\pi^2 i} \frac{2 m k^2 q_\mu}{[m^2 - (k - q)^2 - i0][m^2 - k^2 - i0]} \frac{1}{[v \cdot k + i0]^2} \right\}. \end{aligned} \quad (3.5)$$

Following the prescription of applied in our effective quark model [6,10–16] (see also Ref. [17]) for the calculation of constituent quark loop diagrams we have held only the leading divergent contributions dropping out the contributions independent of the cut-off. This prescription is nothing more than a trivial phenomenological description of quark confinement. Since the appearance of free quarks in the intermediate states of low-energy hadron interactions described by the constituent quark loop diagrams with constant constituent quark masses is related to the imaginary parts of these diagrams, the trivial prohibition of such an appearance is to keeping the divergent contributions of quark loop diagrams related to the real parts of quark diagrams. Moreover, the divergent parts of quark diagrams, being polynomials in powers of the momenta of interacting hadrons, provide local effective low-energy interactions of hadrons in agreement with effective Chiral Lagrangian approach. In terms of the cut-off and the constituent quark mass this procedure restores

fully all effective coupling constants of low-energy interactions of low-lying mesons represented at the hadron level by effective Chiral Lagrangians [10].

The integral containing k_μ in the integrand has the following Lorentz structure: $A(v \cdot q, q^2) v_\mu + B(v \cdot q, q^2) q_\mu$. The contribution of the term linear in v_μ is of order $O(1/M_c^2)$ due to $Q_\mu e^\mu(Q) = 0$ and should be dropped out. Therefore, only the terms linear in q_μ should be held and give the contributions leading in chiral expansion. The integration over k gives one

$$\begin{aligned} \int \frac{d^4 k}{\pi^2 i} \frac{4(m^2 - k^2)k_\mu + 2k^2 q_\mu}{[m^2 - (k - q)^2 - i0][m^2 - k^2 - i0]} \frac{1}{[v \cdot k + i0]} &= -\frac{2}{3} \bar{v}' q_\mu, \\ \int \frac{d^4 k}{\pi^2 i} \frac{2mk^2 q_\mu}{[m^2 - (k - q)^2 - i0][m^2 - k^2 - i0]} \frac{1}{[v \cdot k + i0]^2} &= \\ &= -8m \ell n \left(\frac{\bar{v}'}{4m} \right) q_\mu. \end{aligned} \quad (3.6)$$

For the computation of the integrals in Eq.(3.6) we have used the integrals

$$\begin{aligned} \int \frac{d^4 k}{\pi^2 i} \frac{k_\mu k_\nu}{[m^2 - k^2 - i0]^2} \frac{1}{[v \cdot k + i0]} &= \frac{2}{3} \bar{v}' (g_{\mu\nu} - v_\mu v_\nu), \\ \int \frac{d^4 k}{\pi^2 i} \frac{1}{[m^2 - k^2 - i0]} \frac{1}{[v \cdot k + i0]} &= -\bar{v}', \\ \int \frac{d^4 k}{\pi^2 i} \frac{1}{[m^2 - k^2 - i0]} \frac{1}{[v \cdot k + i0]^2} &= 4 \ell n \left(\frac{\bar{v}'}{4m} \right). \end{aligned} \quad (3.7)$$

We should accentuate that no logarithmically divergent contributions have been omitted in the linearly divergent momentum integrals. Indeed, for example, one can show that

$$\begin{aligned} \int \frac{d^4 k}{\pi^2 i} \frac{1}{[m^2 - k^2 - i0]} \frac{1}{[v \cdot k + i0]} &= -\frac{1}{\pi} \int \frac{d^3 k}{\vec{k}^2 + m^2} = \\ &= -4\Lambda + 4m \operatorname{arctg} \frac{\Lambda}{m} \rightarrow -\bar{v}' + 2\pi m. \end{aligned} \quad (3.8)$$

Thus, following our phenomenological description of confinement, demanding to drop out the terms independent of the cut-off, and collecting all divergent terms of order $O(1/M_c)$ we obtain

$$\begin{aligned} M^{(1)}(D^{*+}(Q) \rightarrow D^0(p)\pi^+(q)) &= \\ &= \sqrt{2} g_{\pi^+} \frac{\sqrt{M_D M_{D^*}}}{\bar{v}'} \left[\frac{m}{M_c} \ell n \left(\frac{\bar{v}'}{4m} \right) - \frac{\bar{v}'}{12M_c} \right] q \cdot e(Q). \end{aligned} \quad (3.9)$$

This gives the coupling constant $g_{D^*D\pi}$ to next-to-leading order in large M_c expansion

$$\begin{aligned} g_{D^*D\pi} &= \sqrt{2} g_{\pi^+} \frac{\sqrt{M_D M_{D^*}}}{\bar{v}'} \left\{ \ell n \left(\frac{\bar{v}'}{4m} \right) + \left[\frac{m}{M_c} \ell n \left(\frac{\bar{v}'}{4m} \right) - \frac{\bar{v}'}{12M_c} \right] \right\} \\ &= 5.2 \times (1 + 7.3 \times 10^{-3}) = 5.24. \end{aligned} \quad (3.10)$$

Thus, we have found that $1/M_c$ corrections to the constant $g_{D^*D\pi}$ make up less than 1% and can be neglected. The accounting of the logarithmically divergent contributions

together with the linearly ones does not go beyond the scope of the approach. Indeed, for the correct calculation of the kinetic terms of the low-lying scalar mesons related to the kinetic terms of the low-lying pseudoscalar ones via chiral transformations one needs to take into account both quadratically and logarithmically divergent contributions [10].

For the computation of the coupling constant $g_{\bar{B}^*\bar{B}\pi}$ we, following the procedure expounded above, get

$$\begin{aligned} g_{\bar{B}^*\bar{B}\pi} &= \sqrt{2} g_{\pi^+} \frac{\sqrt{M_{\bar{B}} M_{\bar{B}^*}}}{\bar{v}'} \left\{ \ln\left(\frac{\bar{v}'}{4m}\right) + \left[\frac{m}{M_b} \ln\left(\frac{\bar{v}'}{4m}\right) - \frac{\bar{v}'}{12M_b} \right] \right\} \\ &\simeq 14.3, \end{aligned} \quad (3.11)$$

where the numerical value is obtained at $M_{\bar{B}^*} = M_{\bar{B}} = M_b = 5.3 \text{ GeV}$ [1]. Thus, we predict negligibly small next-to-leading corrections in heavy M_b expansion to the coupling constant $g_{\bar{B}^*\bar{B}\pi} = 14.3$.

4 Conclusion

When applying the effective quark model with chiral $U(3) \times U(3)$ symmetry incorporating HQEF and $(\text{CHPT})_q$ to the computation the coupling constants $g_{D^*D\pi}$ and $g_{\bar{B}^*\bar{B}\pi}$ to next-to-leading order in heavy-quark mass expansion, we have computed at leading order in large N expansion and in the chiral limit the numerical values $g_{D^*D\pi} = 5.2$ and $g_{\bar{B}^*\bar{B}\pi} = 14.3$ compared reasonably well with the results obtained by Colangelo et al. [2] within QCD sum rules approach and by Dosch and Narison [3] within *full* QCD. Then, we have shown that next-to-leading corrections in large heavy-quark mass expansion give negligible contributions and the main values of the coupling constants are induced at leading order. Our results as well as those predicted by Colangelo et al. [2] and Dosch and Narison [3] are a factor of 2 less compared with those obtained by Belyaev et al.[4]. Therefore, in comparison with Ref.[4] we predict much less contribution of vector mesons D^* and $\bar{B}^*$ to the form factors of the semileptonic decays $D \rightarrow \pi \ell \nu_\ell$ and $\bar{B} \rightarrow \pi \ell \nu_\ell$. The former is very important for the understanding of a mechanism of the semileptonic decays $D \rightarrow \pi \ell \nu_\ell$ and $\bar{B} \rightarrow \pi \ell \nu_\ell$. It should be stressed that our numerical results contain a theoretical uncertainty $\pm \Delta$, the rough estimate of which gives $\Delta \simeq 50\%$ [6]. However for all applications of HQET and $(\text{CHPT})_q$ [6,11–16] the resultant agreements between theoretical and experimental data turn out much better.

5 Acknowledgement

The authors would like to thank President Professor Abdus Salam, the International Atomic Energy Agency and UNESCO for hospitality at the International Centre for Theoretical Physics, Trieste, where the main part of this paper has been carried out.

References

- [1] Partical Data Group, *Phys. Rev. D*, 54 (1996) 1, Part I.
- [2] Colangelo P., Nardulli G., Deandrea A., Di Bartolomeo N., Gatto R. and Feruglio F., *Phys. Lett. B*, 339 (1994) 151.
- [3] Dosch H. G. and Narison S., *Phys. Lett. B* 368 (1996) 163.
- [4] Belyaev V. M., Braun V. M., Khodjamirian A. and Rückl R., *Phys. Rev. D*, 51 (1995) 6167 and references therein.
- [5] Shufman M. A., Vainshtein A. I. and Zakharov V. I., *Nucl. Phys. B*, 147 (1979) 385, 448.
- [6] Ivanov A. N. and Troitskaya N. I. , *Phys. Lett. B*, 345 (1995) 175.
- [7] Eichten E. and Feinberg F. L., *Phys. Rev. D*, 23 (1981) 2724; Eichten E., *Nucl. Phys. B*, 4 (Proc.Suppl.) (1988) 70; Voloshin M. B. and Shifman M. A., *Sov. J. Nucl. Phys.*, 45 (1987) 292;
- [8] Politzer H. D. and Wise M., *Phys. Lett. B*, 206 (1988); *ibid. B*, 208 (1988) 504.
- [9] Georgi H., *Phys. Lett. B*, 240 (1990) 447.
- [10] Ivanov A. N., Nagy M. and Troitskaya N. I., *Int. J. Mod. Phys. A*, 7 (1992) 7305; Ivanov A. N. , *Int. J. Mod. Phys. A*, 8 (1993) 853; Ivanov A. N., Troitskaya N. I. and Nagy M., *Int. J. Mod. Phys. A*, 8 (1993) 2027; 3425; Ivanov A. N., Troitskaya N. I. and Nagy M., *Phys. Lett. B*, 308 (1993) 111; Ivanov A. N. and Troitskaya N. I., *Nuovo Cim. A*, 108 (1995) 555.
- [11] Hussain F., Ivanov A. N. and Troitskaya N. I., *Phys. Lett. B*, 329 (1994) 98; *ibid. B*, 334 (1994) E450; Ivanov A. N., Troitskaya N. I. and Nagy M., *Phys. Lett. B*, 339 (1994) 167.
- [12] Hussain F., Ivanov A. N. and Troitskaya N. I., *Phys. Lett. B*, 348 (1995) 609; *ibid. B*, 369 (1996) 351.
- [13] Ivanov A. N. and Troitskaya N. I., *Phys. Lett. B*, 342 (1995) 323.
- [14] Ivanov A. N. and Troitskaya N. I., *Phys. Lett. B*, 390 (1997) 341.
- [15] Ivanov A. N. and Troitskaya N. I., *Phys. Lett. B*, 394 (1997) 195.
- [16] Ivanov A. N. and Troitskaya N. I., *Nuovo Cim. A*, 110 (1997) 65.
- [17] Bardeen W. A. and Hill C. T., *Phys. Rev. D*, 49 (1994) 409.